


# Spotlight on MAYZENT<sup>®</sup>: Viewing Guide

A HELPFUL RESOURCE  
AS YOU WATCH



Use this guide as you watch the videos to keep track of your thoughts, your feelings, and the things you just don't want to forget! Hang on to it, and use any notes you take to help make your next conversation with your health care provider more productive.

## ON-DEMAND VIDEO GUIDE CONTENTS

**Video 1:** Understanding How RMS Changes Over Time

---

**Video 2:** The Science Behind How MAYZENT Works

---

**Video 3:** The Effectiveness of MAYZENT

---

**Video 4:** MAYZENT Safety

---

**Video 5:** Alongside<sup>™</sup> MAYZENT and Taking the Next Step

---

Additional resources

RMS=relapsing multiple sclerosis.

### What is MAYZENT<sup>®</sup> (siponimod) tablets?

MAYZENT is a prescription medicine that is used to treat relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults. It is not known if MAYZENT is safe and effective in children.

### IMPORTANT SAFETY INFORMATION

#### Do not take MAYZENT if you:

- have a CYP2C9\*3/\*3 genotype. Before starting treatment with MAYZENT, your CYP2C9 genotype should be determined by your health care provider. Ask your health care provider if you are not sure.
- have had a heart attack, chest pain called unstable angina, stroke or mini-stroke (transient ischemic attack or TIA), or certain types of heart failure in the last 6 months
- have certain types of heart block or irregular or abnormal heartbeat (arrhythmia), unless you have a pacemaker

Please see additional Important Safety Information throughout and full [Prescribing Information](#), including [Medication Guide](#). **1**

## VIDEO 1: UNDERSTANDING HOW RMS CHANGES OVER TIME

---

1. Is my RMS feeling different recently?

---

2. Am I experiencing symptoms that are new or worsening to me?

---

3. Am I having trouble doing things that once came more easily to me? Take a look at the key areas for determining disability below, and make a note of any that you are experiencing.



Assistance required  
to walk



Trouble with memory  
or concentration



Weakness or problems  
moving limbs



Vision problems



Problems with  
coordination



Difficulty with speech  
or swallowing



Bowel and  
bladder issues



Numbness or loss  
of sensation

## VIDEO 2: THE SCIENCE BEHIND HOW MAYZENT<sup>®</sup> WORKS

---

1. What would taking a once-daily oral treatment change in my life?

---

2. Is there anything I'd like to ask my doctor about how MAYZENT may work?

---

3. Have I learned anything new about how MAYZENT is thought to work?

---

### IMPORTANT SAFETY INFORMATION *(cont)*

**MAYZENT may cause serious side effects, including:**

**1. Slow heart rate (bradycardia or bradyarrhythmia) when you start taking MAYZENT.** MAYZENT can cause your heart rate to slow down, especially after you take your first dose. You should have a test to check the electrical activity of your heart called an electrocardiogram (ECG) before you take your first dose of MAYZENT.

### **VIDEO 3: THE EFFECTIVENESS OF MAYZENT<sup>®</sup>**

---

1. What are some of the treatment goals that I can set for myself?

---

2. Do I know my Expanded Disability Status Scale (EDSS) score? Why is my EDSS score important?

---

3. What could more time without disability progression mean for me?

---

### **VIDEO 4: MAYZENT SAFETY**

---

1. What are the most common side effects with MAYZENT?

---

2. Is there anything I'd like to tell my doctor before starting treatment with MAYZENT?

---

3. Is there anything else about safety I want to discuss with my doctor?

---

### **VIDEO 5: ALONGSIDE<sup>™</sup> MAYZENT AND TAKING THE NEXT STEP**

---

1. Which of the Alongside MAYZENT support offerings am I interested in?

---

2. How can I find out if I'm eligible for financial support with my treatment?

---

3. Do I have any other questions about starting treatment with MAYZENT?

---

#### **IMPORTANT SAFETY INFORMATION** *(cont)*

**1. Slow heart rate (bradycardia or bradyarrhythmia) when you start taking MAYZENT (cont):** During the initial up dosing period (4 days for the 1-mg daily dose or 5 days for the 2-mg daily dose), if you miss 1 or more doses of MAYZENT, you need to restart the up dosing. Call your health care provider if you miss a dose of MAYZENT.

**2. Infections.** MAYZENT can increase your risk of serious infections that can be life-threatening and cause death.

## ADDITIONAL RESOURCES

---

For more information on all things MAYZENT<sup>®</sup>, use the helpful resources listed below. To find all of these resources online, visit <https://www.mayzent.com/resources>.

### **MAYZENT DISCUSSION GUIDE**

Answer a few questions to generate a personalized discussion guide that can help your doctor capture a picture of your condition.

### **DISCOVER MAYZENT BROCHURE**

Facts you should know about MAYZENT all in one downloadable brochure.

### **READY TO START MAYZENT BROCHURE**

A step-by-step guide to begin treatment with MAYZENT.

### **THE CAREGIVER'S GUIDEBOOK**

A downloadable book that can help you navigate a loved one's journey with multiple sclerosis (MS). In it you'll find information, strategies, and support—a valuable read for an MS care partner.

[DOWNLOAD](#)



### **START FORM**

Download a Start Form today, fill out your information, and bring it to your next doctor's appointment. This can help you get a head start on starting MAYZENT.

### **IMPORTANT SAFETY INFORMATION** *(cont)*

**2. Infections (cont):** MAYZENT lowers the number of white blood cells (lymphocytes) in your blood. This will usually go back to normal within 3 to 4 weeks of stopping treatment. Your health care provider should review a recent blood test of your white blood cells before you start taking MAYZENT.

### IMPORTANT SAFETY INFORMATION (cont)

**2. Infections (cont):** Call your health care provider right away if you have any of these symptoms of an infection during treatment with MAYZENT and for 3 to 4 weeks after your last dose of MAYZENT:

- fever
- tiredness
- body aches
- chills
- nausea
- vomiting
- headache with fever, neck stiffness, sensitivity to light, nausea, confusion (these may be symptoms of meningitis, an infection of the lining around your brain and spine)

**3. A problem with your vision called macular edema.** Macular edema can cause some of the same vision symptoms as a multiple sclerosis (MS) attack (optic neuritis). You may not notice any symptoms with macular edema. If macular edema happens, it usually starts in the first 1 to 4 months after you start taking MAYZENT. Your health care provider should test your vision before you start taking MAYZENT and any time you notice vision changes during treatment with MAYZENT. Your risk of macular edema is higher if you have diabetes or have had an inflammation of your eye called uveitis.

Call your health care provider right away if you have any of the following: blurriness or shadows in the center of your vision, a blind spot in the center of your vision, sensitivity to light, or unusually colored (tinted) vision.

**Before taking MAYZENT, tell your health care provider about all of your medical conditions, including if you:**

- have an irregular or abnormal heartbeat
- have a history of stroke or other diseases related to blood vessels in the brain
- have breathing problems, including during your sleep
- have a fever or infection, or you are unable to fight infections due to a disease or are taking medicines that lower your immune system. Tell your health care provider if you have had chickenpox or have received the vaccine for chickenpox. Your health care provider may do a blood test for chickenpox virus. You may need to get the full course of vaccine for chickenpox and then wait 1 month before you start taking MAYZENT.
- have slow heart rate
- have liver problems

- have diabetes
- have eye problems, especially an inflammation of the eye called uveitis
- had or now have a type of skin cancer called basal cell carcinoma (BCC), melanoma, or squamous cell carcinoma
- have high blood pressure
- are pregnant or plan to become pregnant. MAYZENT may harm your unborn baby. Talk to your health care provider right away if you become pregnant while taking MAYZENT or if you become pregnant within 10 days after you stop taking MAYZENT.
  - If you are a woman who can become pregnant, you should use effective birth control during your treatment with MAYZENT and for at least 10 days after you stop taking MAYZENT.
- are breastfeeding or plan to breastfeed. It is not known if MAYZENT passes into your breast milk. Talk to your health care provider about the best way to feed your baby if you take MAYZENT.

**Tell your health care provider about all the medicines you take, including** prescription medicines, over-the-counter medicines, vitamins, and herbal supplements. Especially tell your health care provider if you take medicines to control your heart rhythm (anti-arrhythmics), or blood pressure (antihypertensives), or heart beat (such as calcium channel blockers or beta-blockers); take medicines that affect your immune system, such as beta-interferon or glatiramer acetate, or any of these medicines that you took in the past.

Tell your health care provider if you have recently received a live vaccine. You should avoid receiving **live** vaccines during treatment with MAYZENT. MAYZENT should be stopped 1 week before and for 4 weeks after receiving a live vaccine. If you receive a live vaccine, you may get the infection the vaccine was meant to prevent. Vaccines may not work as well when given during treatment with MAYZENT.

**MAYZENT may cause possible side effects, including:**

- **increased blood pressure.** Your health care provider should check your blood pressure during treatment with MAYZENT.

**IMPORTANT SAFETY INFORMATION** (cont)

**MAYZENT may cause possible side effects, including (cont):**

- **liver problems.** MAYZENT may cause liver problems. Your health care provider should do blood tests to check your liver before you start taking MAYZENT. Call your health care provider right away if you have any of the following symptoms of liver problems:
  - nausea
  - vomiting
  - stomach pain
  - tiredness
  - loss of appetite
  - your skin or the whites of your eyes turn yellow
  - dark urine
- **breathing problems.** Some people who take MAYZENT have shortness of breath. Call your health care provider right away if you have new or worsening breathing problems.
- **swelling and narrowing of the blood vessels in your brain.** A condition called PRES (Posterior Reversible Encephalopathy Syndrome) has happened with drugs in the same class. Symptoms of PRES usually get better when you stop taking MAYZENT. However, if left untreated, it may lead to a stroke. Call your health care provider right away if you have any of the following symptoms: sudden severe headache, sudden confusion, sudden loss of vision or other changes in vision, or seizure.
- **severe worsening of multiple sclerosis after stopping MAYZENT.** When MAYZENT is stopped, symptoms of MS may return and become worse compared to before or during treatment. Always talk to your doctor before you stop taking MAYZENT for any reason. Tell your health care provider if you have worsening symptoms of MS after stopping MAYZENT.
- **a type of skin cancer called basal cell carcinoma (BCC), melanoma, and squamous cell carcinoma.** Tell your doctor if you have any changes in the appearance of your skin, including changes in a mole, a new darkened area on your skin, a sore that does not heal, or growths on your skin, such as a bump that may be shiny, pearly white, skin-colored, or pink. Your doctor should check your skin for any changes during treatment with MAYZENT. Limit the amount of time you spend in sunlight and ultraviolet (UV) light. Wear protective clothing and use a sunscreen with a high sun protection factor.

**The most common side effects of MAYZENT include:**

headache, high blood pressure (hypertension), and abnormal liver tests.

These are not all of the possible side effects of MAYZENT. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

**Please see full [Prescribing Information](#), including [Medication Guide](#).**

MAYZENT and the MAYZENT logo are registered trademarks of Novartis AG. Alongside MAYZENT is a trademark of Novartis AG. All other trademarks are the property of their respective owners.